Cabrera, A. R., et al. (2015). "Initial recommendations for higher-tier risk assessment protocols for bumble bees, Bombus spp.(Hymenoptera: Apidae)." <u>Integrated</u> environmental assessment and management.

Cutler, G. C. and C. D. Scott-Dupree (2014). "A field study examining the effects of exposure to neonicotinoid seed-treated corn on commercial bumble bee colonies." <u>Ecotoxicology</u> **23**(9): 1755-1763.

Dively, G. P., et al. (2015). "Assessment of chronic sublethal effects of imidacloprid on honey bee colony health." <u>PloS one</u> **10**(3): e0118748.

Doublet, V., et al. (2015). "Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle." <u>Environmental microbiology</u> **17**(4): 969-983.

Frazier, M. T., et al. (2015). "Assessing Honey Bee (Hymenoptera: Apidae) Foraging Populations and the Potential Impact of Pesticides on Eight US Crops." <u>Journal of Economic Entomology</u>: tov195.

Garbuzov, M., et al. (2015). "Honey bee dance decoding and pollen-load analysis show limited foraging on spring-flowering oilseed rape, a potential source of neonicotinoid contamination." <u>Agriculture, Ecosystems & Environment</u> **203**: 62-68.

Jin, N., et al. (2015). "The neonicotinoid clothianidin interferes with navigation of the solitary bee Osmia cornuta in a laboratory test." <u>The Journal of experimental biology</u>: jeb. 123612.

Kessler, S. C., et al. (2015). "Bees prefer foods containing neonicotinoid pesticides." <u>Nature</u>.

Lundin, O., et al. (2015). "Neonicotinoid Insecticides and Their Impacts on Bees: A Systematic Review of Research Approaches and Identification of Knowledge Gaps." <u>PloSone</u> **10**(8): e0136928.

Raine, N. E. and R. J. Gill (2015). "Ecology: Tasteless pesticides affect bees in the field." Nature.

Rundlöf, M., et al. (2015). "Seed coating with a neonicotinoid insecticide negatively affects wild bees." <u>Nature</u> **521**(7550): 77-80.

Samson-Robert, O., et al. (2014). "Neonicotinoid-contaminated puddles of water represent a risk of intoxication for honey bees." <u>PloS one</u> **9**(12): e108443.

Sandrock, C., et al. (2014). "Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success." <u>Agricultural and Forest Entomology</u> **16**(2): 119-128.

Schaafsma, A., et al. (2015). "Neonicotinoid Insecticide Residues in Surface Water and Soil Associated with Commercial Maize (Corn) Fields in Southwestern Ontario." <u>PloS one</u> **10**(2): e0118139.

Scholer, J. and V. Krischik (2014). "Chronic exposure of imidacloprid and clothianidin reduce queen survival, foraging, and nectar storing in colonies of Bombus impatiens." PloS one **9**(3): e91573.

Staveley, J. P., et al. (2014). "A causal analysis of observed declines in managed honey bees (Apis mellifera)." <u>Human and Ecological Risk Assessment: An International Journal</u> **20**(2): 566-591.

Van der Sluijs, J., et al. (2015). "Conclusions of the Worldwide Integrated Assessment on the risks of neonicotinoids and fipronil to biodiversity and ecosystem functioning." <u>Environmental Science and Pollution Research</u> **22**(1): 148-154.

van der Zee, R., et al. (2015). "An Observational Study of Honey Bee Colony Winter Losses and Their Association with Varroa destructor, Neonicotinoids and Other Risk Factors." <u>PloS one</u> **10**(7).